



BOMO™

1.0-0-0.8

Foliar Nutritional Liquid to enhance plant growth

Guaranteed Analysis

Total Nitrogen(N) 1.0%
 0.93% Urea nitrogen
 0.07% ammoniacal nitrogen
 Soluble Potash (K₂O) 0.8%
 Boron(B) 0.20%
 Molybdenum(MO) 0.90%

Derived from Urea, potassium sulfate, Ammonium molybdate tetrahydrate
 (NH₄)₂MO₇O₂₄ · 4H₂O and Boric acid.

WARNING:

This product contains Boron, application to any crops other than those recommended may result in serious injury. The application of fertilizing material containing molybdenum (MO) may result in forage crops containing levels of molybdenum (MO) which are toxic to ruminants.

Keep Out Of Reach Of Children

CAUTION: Store in Cool and dry Conditions

Precautionary Statement:

Before using this product, read all precautions, directions for use, conditions of sale-limited warranty and limitations of liability and remedies.



NCA Biotech, Inc
 3406 Pomona Blvd
 Pomona, CA 91768

Email: info@NCABiotech.com
 Tel: 1 (909) 348-5133 · Fax: 1(909) 348-5135

Net Contents: 1 U.S. Gallon (3.785 L)
 Net Weight.: 8.85 lbs.

BOMO™ is a unique and superior foliar nutritional liquid especially effective to enhance plant physiological activities and plant growth of various food crops. **BOMO™** is specially designed to meet the nutritional requirement of various crops listed below.

Directions for Use:

Crop	Application Rate/ Fl. Oz. product per acre	Application method/Timing	Remarks
Legume vegetables(Succulent or Dried), such as, beans, peas, soybean, broad bean, lentil, & others	5 fl. Oz product	1st application applies at 5-10 % bloom. Repeat applications at 7-10 days interval until fruiting stages	5-10 gals spray volume per acre
Pome/Stone fruits, such as apple, apricot, cherry, plum, plumcot and peach	64 fl. oz product	1st application at 7-14 days prior to flowering. 2nd application at full bloom. 3rd application at small fruit stage.	100-200 gals spray volume per acre
Cotton	10 fl.oz product by ground. 2-5 fl. Oz product by air	1st application at early flowering stage and repeat every 10 days until harvest if required	10-20 gals spray volume per acre by ground. 2-8 gals spray volume per acre by air
Nut crops, such as almond, walnuts and others	62 fl oz product	1st application at early flowering stage. Repeat applications at 7-10 days interval until harvest as required	100-200 gals spray volume per acre
Canola	5 fl. Oz product	1st application applies at 5-10 % bloom. Repeat applications at 7-10 days interval until fruiting stages	5-20 gals spray volume per acre
Other crops	1:200 dilution rate	1st application applies at 5-10 % bloom. Repeat applications at 7-10 days.	

Do not tank mix with any agrichemical products that are alkaline in nature. Make sure the final spray mix is below pH 7.0.

CONTAINER DISPOSAL: Do not re-use empty container, Rinse thoroughly before discarding according Federal and State laws

LIMITED WARRANTY AND DISCLAIMER

NOTICE: NCA BIOTECH, INC., warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. NCA BIOTECH, INC., makes no other express or implied warranty of fitness or merchantability. In no case shall NCA BIOTECH, INC., or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. NCA Biotech, Inc, and Seller offer this product and the Buyer and user accept it, subject to the foregoing **Limited Warranty and Disclaimer** which may be varied only by agreement in writing signed by a duly authorized representative of NCA BIOTECH, INC

BOMO™ is a trade mark of NCA Biotech, Inc

NCA Biotech, Inc
 3406 Pomona Blvd
 Pomona, CA 91768
 Email: info@NCABiotech.com
 Tel: 1 (909) 348-5133 · Fax: 1(909) 348-5135